



Rad54 rabbit pAb

Cat No.:ES7755

For research use only


Overview

Product Name	Rad54 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human RAD54L. AA range:221-270
Specificity	Rad54 Polyclonal Antibody detects endogenous levels of Rad54 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	DNA repair and recombination protein RAD54-like
Gene Name	RAD54L
Cellular localization	Nucleus .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	85kD
Human Gene ID	8438
Human Swiss-Prot Number	Q92698
Alternative Names	RAD54L; RAD54A; DNA repair and recombination protein RAD54-like; RAD54 homolog; hHR54; hRAD54
Background	The protein encoded by this gene belongs to the DEAD-like helicase superfamily, and shares similarity with <i>Saccharomyces cerevisiae</i> Rad54, a protein






known to be involved in the homologous recombination and repair of DNA. This protein has been shown to play a role in homologous recombination related repair of DNA double-strand breaks. The binding of this protein to double-strand DNA induces a DNA topological change, which is thought to facilitate homologous DNA pairing, and stimulate DNA recombination. Alternative splicing results in multiple transcript variants encoding the same protein.[provided by RefSeq, Dec 2008],



Western blot analysis of lysates from HeLa cells, Jurkat cells, and 293 cells, using RAD54 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

