



COX4I2 rabbit pAb

Cat No.:ES7777

For research use only


Overview

Product Name	COX4I2 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human COX42. AA range:31-80
Specificity	COX4I2 Polyclonal Antibody detects endogenous levels of COX4I2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Cytochrome c oxidase subunit 4 isoform 2 mitochondrial
Gene Name	COX4I2
Cellular localization	Mitochondrion inner membrane ; Single-pass membrane protein .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	20kD
Human Gene ID	84701
Human Swiss-Prot Number	Q96KJ9
Alternative Names	COX4I2; COX4L2; Cytochrome c oxidase subunit 4 isoform 2; mitochondrial; Cytochrome c oxidase subunit IV isoform 2; COX IV-2
Background	Cytochrome c oxidase (COX), the terminal enzyme of







the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. It is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may be involved in the regulation and assembly of the complex. This nuclear gene encodes isoform 2 of subunit IV. Isoform 1 of subunit IV is encoded by a different gene, however, the two genes show a similar structural organization. Subunit IV is the largest nuclear encoded subunit which plays a pivotal role in COX regulation. [provided by RefSeq, Jul 2008],




Western Blot analysis of K562 insulin 0.01U/ml 15' cells using COX4I2 Polyclonal Antibody

Immunofluorescence analysis of COS7 cells, using COX42 Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from K562 cells, treated with insulin 0.01U/ml 15', using COX42 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

