

DD3 rabbit pAb

Cat No.:ES7844

For research use only

Overview

Product Name	DD3 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not
	yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human AKR1C3. AA
	range:191-240
Specificity	DD3 Polyclonal Antibody detects endogenous levels
	of DD3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Aldo-keto reductase family 1 member C3
Gene Name	AKR1C3
Cellular localization	Cytoplasm .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	37kD
Human Gene ID	8644
Human Swiss-Prot Number	P42330
Alternative Names	AKR1C3; DDH1; HSD17B5; KIAA0119; PGFS;
	Aldo-keto reductase family 1 member C3;
	17-beta-hydroxysteroid dehydrogenase type 5;
	17-beta-HSD 5; 3-alpha-HSD type II; brain;
	3-alpha-hydroxysteroid dehydrogenase type 2;
	3-alpha-HSD type 2; Chlordec
Background	This gene encodes a member of the aldo/keto
	reductase superfamily, which consists of more than



+86-27-59760950


ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols by utilizing NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme catalyzes the reduction of prostaglandin (PG) D2, PGH2 and phenanthrenequinone (PQ), and the oxidation of 9alpha,11beta-PGF2 to PGD2. It may play an important role in the pathogenesis of allergic diseases such as asthma, and may also have a role in controlling cell growth and/or differentiation. This gene shares high sequence identity with three other gene members and is clustered with those three genes at chromosome 10p15-p14. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2011],



Western blot analysis of lysates from HeLa and COLO cells, using AKR1C3 Antibody. The lane on the right is blocked with the synthesized peptide.



ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C