

## c-FLIP rabbit pAb

## Cat No.:ES7886

For research use only

## Overview

Product Name	c-FLIP rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human in ternal CFLAR. AA
	range:181-230
Specificity	c-FLIP Polyclonal Antibody detects endogenous
	levels of c-FLIP protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	CASP8 and FADD-like apoptosis regulator
Gene Name	CFLAR
Cellular localization	cytoplasm,cytosol,death-inducing signaling
	complex,CD95 death-inducing signaling
	complex,membrane raft,ripoptosome,
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	55kD
Human Gene ID	8837
Human Swiss-Prot Number	015519
Alternative Names	CFLAR; CASH; CASP8AP1; CLARP; MRIT; CASP8 and
	FADD-like apoptosis regulator; Caspase homolog;
	CASH; Caspase-eight-related protein; Casper;
	Caspase-like apoptosis regulatory protein; CLARP;
	Cellular FLICE-like inhibitory protein; c-FLIP; FAD



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Background

The protein encoded by this gene is a regulator of apoptosis and is structurally similar to caspase-8. However, the encoded protein lacks caspase activity and appears to be itself cleaved into two peptides by caspase-8. Several transcript variants encoding different isoforms have been found for this gene, and partial evidence for several more variants exists. [provided by RefSeq, Feb 2011],

Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C