

SFRS11 rabbit pAb

Cat No.: ES8014

For research use only

Overview

Product Name SFRS11 rabbit pAb

Host species Rabbit

Applications WB;ELISA;IHC Species Cross-Reactivity Human;Mouse

Recommended dilutions WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000

Immunogen The antiserum was produced against synthesized peptide derived from human SFRS11. AA

range:211-260

Specificity SFRS11 Polyclonal Antibody detects endogenous

levels of SFRS11 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Serine/arginine-rich splicing factor 11

Gene Name SRSF11

Cellular localization Nucleus. Colocalizes with spliceosome components.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 54kD
Human Gene ID 9295
Human Swiss-Prot Number Q05519

Alternative Names SRSF11; SFRS11; Serine/arginine-rich splicing factor

11; Arginine-rich 54 kDa nuclear protein; p54;

Splicing factor; arginine/serine-rich 11

Background This gene encodes 54-kD nuclear protein that

contains an arginine/serine-rich region similar to segments found in pre-mRNA splicing factors. Although the function of this protein is not yet known, structure and immunolocalization data suggest that it may play a role in pre-mRNA



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



processing. Alternative splicing results in multiple transcript variants encoding different proteins. In addition, a pseudogene of this gene has been found on chromosome 12.[provided by RefSeq, Sep 2010],

Western blot analysis of lysates from LOVO and RAW264.7 cells, using SFRS11 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

