



# RIP140 (Acetyl Lys158) rabbit pAb

Cat No.:ES8434

For research use only


## Overview

<b>Product Name</b>	RIP140 (Acetyl Lys158) rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;ELISA;IHC
<b>Species Cross-Reactivity</b>	Human;Mouse;Rat
<b>Recommended dilutions</b>	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
<b>Immunogen</b>	Synthesized acetyl-peptide derived from human RIP140 around the acetylation site of K158.
<b>Specificity</b>	Acetyl-RIP140 (K158) Polyclonal Antibody detects endogenous levels of RIP140 around the acetylation site of K158 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	Nuclear receptor-interacting protein 1
<b>Gene Name</b>	NRIP1
<b>Cellular localization</b>	Nucleus . Localized to discrete foci and redistributes to larger nuclear domains upon binding to ligand-bound NR3C1.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	
<b>Human Gene ID</b>	8204
<b>Human Swiss-Prot Number</b>	P48552
<b>Alternative Names</b>	NRIP1; Nuclear receptor-interacting protein 1; Nuclear factor RIP140; Receptor-interacting protein 140
<b>Background</b>	Nuclear receptor interacting protein 1 (NRIP1) is a nuclear protein that specifically interacts with the hormone-dependent activation domain AF2 of nuclear receptors. Also known as RIP140, this





protein modulates transcriptional activity of the estrogen receptor. [provided by RefSeq, Jul 2008],



Western blot analysis of 293T using Acetyl-RIP140 (K158) antibody. Antibody was diluted at 1:500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

