

## RL18A rabbit pAb

Cat No.:ES9321

For research use only

## Overview

Product Name RL18A rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

**Immunogen** Synthesized peptide derived from human protein .

at AA range: 100-180

**Specificity** RL18A Polyclonal Antibody detects endogenous

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

**Protein Name** 60S ribosomal protein L18a

Gene Name RPL18A

**Cellular localization** cytosol,ribosome,membrane,cytosolic large

ribosomal subunit,

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 19kD
Human Gene ID 6142
Human Swiss-Prot Number Q02543

**Alternative Names** 

**Background** Ribosomes, the organelles that catalyze protein


synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a member of the L18AE family of ribosomal proteins that is a component of the 60S subunit. The encoded protein may play a role in viral replication by interacting



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





with the hepatitis C virus internal ribosome entry site (IRES). This gene is co-transcribed with the U68 snoRNA, located within the third intron. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed throughout the genome. [provided by RefSeq, Jul 2012],

Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night



+86-27-59760950