

## PDE1A rabbit pAb

## Cat No.:ES9466

For research use only

## Overview

Product Name	PDE1A rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
<b>Recommended dilutions</b>	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from part region of
	human protein
Specificity	PDE1A Polyclonal Antibody detects endogenous
	levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20 $^\circ\!\mathrm{C}$ . Avoid repeated freeze-thaw cycles.
Protein Name	Calcium/calmodulin-dependent 3',5'-cyclic
	nucleotide phosphodiesterase 1A (Cam-PDE 1A) (EC
	3.1.4.17) (61 kDa Cam-PDE) (hCam-1)
Gene Name	PDE1A
Cellular localization	nucleus,cytosol,neuronal cell body,
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	58kD
Human Gene ID	5136
Human Swiss-Prot Number	P54750
Alternative Names	
Background	Cyclic nucleotide phosphodiesterases (PDEs) play a
	role in signal transduction by regulating intracellular
	cyclic nucleotide concentrations through hydrolysis
	of cAMP and/or cGMP to their respective nucleoside
	5-prime monophosphates. Members of the PDE1
	family, such as PDE1A, are Ca(2+)/calmodulin (see



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

CALM1; MIM 114180)-dependent PDEs (CaM-PDEs)

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



that are activated by calmodulin in the presence of Ca(2+) (Michibata et al., 2001 [PubMed 11342109]; Fidock et al., 2002 [PubMed 11747989]).[supplied by OMIM, Oct 2009],



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C