

NR2C2 rabbit pAb

Cat No.:ES9965

For research use only

Overview

Product Name	NR2C2 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human protein .
-	at AA range: 410-490
Specificity	NR2C2 Polyclonal Antibody detects endogenous
	levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Nuclear receptor subfamily 2 group C member 2
	(Orphan nuclear receptor TAK1) (Orphan nuclear
	receptor TR4) (Testicular receptor 4)
Gene Name	NR2C2 TAK1 TR4
Cellular localization	Nucleus .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	65kD
Human Gene ID	7182
Human Swiss-Prot Number	P49116
Alternative Names	
Background	This gene encodes a protein that belongs to the
	nuclear hormone receptor family. Members of this
	family act as ligand-activated transcription factors
	and function in many biological processes such as
	development, cellular differentiation and
	homeostasis. The activated receptor/ligand complex



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

is translocated to the nucleus where it binds to

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



hormone response elements of target genes. The protein encoded by this gene plays a role in protecting cells from oxidative stress and damage induced by ionizing radiation. The lack of a similar gene in mouse results in growth retardation, severe spinal curvature, subfertility, premature aging, and prostatic intraepithelial neoplasia (PIN) development. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Apr 2014],

Western Blot analysis of mouse-brain rat-brain Hela 293 PC-3 cell lysis. Primary Antibody was diluted at 1:1000. Secondary antibody(catalog#:RS0002 was diluted at 1:10000





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C