

Recombinant Human AOC3 (C-6His)

Catalog #	EPT191
Expression Host	Human Cells
DESCRIPTION	Recombinant Human Membrane Primary Amine
	Oxidase is produced by our Mammalian expression
	system and the target gene encoding Arg28-Asn763 is
	expressed with a 6His tag at the C-terminus.
Accession	Q16853
Synonyms	Membrane primary amine oxidase; Copper amine
	oxidase; HPAO; Semicarbazide-sensitive amine
	oxidase; SSAO; Vascular adhesion protein 1; VAP-1;
	AOC3; VAP1
Mol Mass	82.6 KDa
AP Mol Mass	90-110 KDa, reducing conditions
Purity	Greater than 95% as determined by reducing
	SDS-PAGE.
Endotoxin	Less than 0.1 ng/ μ g (1 EU/ μ g) as determined by LAL
	test.
FORMULATION	Supplied as a 0.2 μ m filtered solution of 20mM



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com 23-2, No.388 Gaoxin 2nd Road,Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.Q



Tris-HCl, 500mM NaCl, pH 8.0.

RECONSTITUTION	
SHIPPING	The product is shipped on dry ice/polar packs.
	Upon receipt, store it immediately at the temperature
	listed below.
STORAGE	Store at \leq -70°C, stable for 6 months after receipt.
	Store at \leq -70 °C, stable for 3 months under sterile
	conditions after opening.
	Please minimize freeze-thaw cycles.
BACKGROUND	Vascular adhesion protein-1(VAP-1) is a copper amine
	oxidase with a topaquinone cofactor.VAP-1 is a type II
	integral membrane protein, but a soluble form of the
	enzyme is present in human serum, and its level
	increases in diabetes and some inflammatory liver
	diseases. VAP-1 catalyzes the oxidative deamination of
	small primary amines such as methylamine,
	benzylamine, and aminoacetone in a reaction that
	produces an aldehyde, ammonia, and H2O2. VAP-1
	vascular expression is regulated at sites of
	inflammation through its release from intracellular
	granules in which the protein is stored. The adhesive
	function of VAP-1 has been demonstrated in studies



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



showing that the protein is important for the adherence of certain lymphocyte subtypes to inflamed endothelial tissues. VAP-1 mediated adhesion is involved in the process of leukocyte extravasation, an important feature of inflammatory responses. VAP-1 is considered to be a therapeutic target for diabetes, oxidative stress, and inflammatory diseases.



SDS-PAGE



+86-27-59760950 ELKbio@ELKbiotech.com

.com www.elkk

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C